

INFORMATION IN THE TABLE BELOW AND IN THE TABLES ON PAGES 3-5 MAY BE USEFUL IN ANSWERING
THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.

DO NOT DETACH FROM BOOK.

PERIODIC TABLE OF THE ELEMENTS

1 H 1.008	2 He 4.00
3 Li 6.94	4 Be 9.01
11 Na 22.99	12 Mg 24.30
19 K 39.10	20 Ca 40.08
21 Sc 44.96	22 Ti 47.90
23 V 50.94	24 Cr 52.00
25 Mn 54.94	26 Fe 55.85
27 Co 58.93	28 Ni 58.69
29 Cu 63.55	30 Zn 65.39
31 Ga 69.72	32 Ge 72.59
33 As 74.92	34 Se 78.96
35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62
39 Y 88.91	40 Zr 91.22
41 Nb 92.91	42 Mo 95.94
43 Tc (98)	44 Ru 101.1
45 Rh 102.91	46 Pd 106.42
47 Ag 107.87	48 Cd 112.41
49 In 114.82	50 Sn 118.71
51 Sb 121.75	52 Te 127.60
53 I 126.91	54 Xe 131.29
55 Cs 132.91	56 Ba 137.33
57 *La 138.91	72 Hf 178.49
73 Ta 180.95	74 W 183.85
75 Re 186.21	76 Os 190.2
77 Ir 192.2	78 Pt 195.08
79 Au 196.97	80 Hg 200.59
81 Tl 204.38	82 Pb 207.2
83 Bi 208.98	84 Po (209)
85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226.02
89 +Ac 227.03	104 Rf (261)
105 Db (262)	106 Sg (266)
107 Bh (264)	108 Hs (277)
109 Mt (268)	110 Ds (271)
111 Rg (272)	

*Lanthanide Series	58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.4	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
	90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (262)

STANDARD REDUCTION POTENTIALS IN AQUEOUS SOLUTION AT 25°C

Half-reaction		$E^\circ(V)$
$F_2(g) + 2e^-$	\rightarrow	2F ⁻ 2.87
$Co^{3+} + e^-$	\rightarrow	Co ²⁺ 1.82
$Au^{3+} + 3e^-$	\rightarrow	Au(s) 1.50
$Cl_2(g) + 2e^-$	\rightarrow	2Cl ⁻ 1.36
$O_2(g) + 4H^+ + 4e^-$	\rightarrow	2H ₂ O(l) 1.23
$Br_2(l) + 2e^-$	\rightarrow	2Br ⁻ 1.07
$2Hg^{2+} + 2e^-$	\rightarrow	Hg ₂ ²⁺ 0.92
$Hg^{2+} + 2e^-$	\rightarrow	Hg(l) 0.85
$Ag^+ + e^-$	\rightarrow	Ag(s) 0.80
$Hg_2^{2+} + 2e^-$	\rightarrow	2Hg(l) 0.79
$Fe^{3+} + e^-$	\rightarrow	Fe ²⁺ 0.77
$I_2(s) + 2e^-$	\rightarrow	2I ⁻ 0.53
$Cu^+ + e^-$	\rightarrow	Cu(s) 0.52
$Cu^{2+} + 2e^-$	\rightarrow	Cu(s) 0.34
$Cu^{2+} + e^-$	\rightarrow	Cu ⁺ 0.15
$Sn^{4+} + 2e^-$	\rightarrow	Sn ²⁺ 0.15
$S(s) + 2H^+ + 2e^-$	\rightarrow	H ₂ S(g) 0.14
$2H^+ + 2e^-$	\rightarrow	H ₂ (g) 0.00
$Pb^{2+} + 2e^-$	\rightarrow	Pb(s) -0.13
$Sn^{2+} + 2e^-$	\rightarrow	Sn(s) -0.14
$Ni^{2+} + 2e^-$	\rightarrow	Ni(s) -0.25
$Co^{2+} + 2e^-$	\rightarrow	Co(s) -0.28
$Cd^{2+} + 2e^-$	\rightarrow	Cd(s) -0.40
$Cr^{3+} + e^-$	\rightarrow	Cr ²⁺ -0.41
$Fe^{2+} + 2e^-$	\rightarrow	Fe(s) -0.44
$Cr^{3+} + 3e^-$	\rightarrow	Cr(s) -0.74
$Zn^{2+} + 2e^-$	\rightarrow	Zn(s) -0.76
$2H_2O(l) + 2e^-$	\rightarrow	H ₂ (g) + 2OH ⁻ -0.83
$Mn^{2+} + 2e^-$	\rightarrow	Mn(s) -1.18
$Al^{3+} + 3e^-$	\rightarrow	Al(s) -1.66
$Be^{2+} + 2e^-$	\rightarrow	Be(s) -1.70
$Mg^{2+} + 2e^-$	\rightarrow	Mg(s) -2.37
$Na^+ + e^-$	\rightarrow	Na(s) -2.71
$Ca^{2+} + 2e^-$	\rightarrow	Ca(s) -2.87
$Sr^{2+} + 2e^-$	\rightarrow	Sr(s) -2.89
$Ba^{2+} + 2e^-$	\rightarrow	Ba(s) -2.90
$Rb^+ + e^-$	\rightarrow	Rb(s) -2.92
$K^+ + e^-$	\rightarrow	K(s) -2.92
$Cs^+ + e^-$	\rightarrow	Cs(s) -2.92
$Li^+ + e^-$	\rightarrow	Li(s) -3.05

ATOMIC STRUCTURE

$$\begin{aligned}E &= h\nu & c &= \lambda\nu \\ \lambda &= \frac{h}{mv} & p &= mv \\ E_n &= \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}\end{aligned}$$

EQUILIBRIUM

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_b = \frac{[\text{OH}^-][\text{HB}^+]}{[\text{B}]}$$

$$\begin{aligned}K_w &= [\text{OH}^-][\text{H}^+] = 1.0 \times 10^{-14} @ 25^\circ\text{C} \\ &= K_a \times K_b\end{aligned}$$

$$\text{pH} = -\log[\text{H}^+], \text{pOH} = -\log[\text{OH}^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{HB}^+]}{[\text{B}]}$$

$$\text{p}K_a = -\log K_a, \text{p}K_b = -\log K_b$$

$$K_p = K_c(RT)^{\Delta n},$$

where Δn = moles product gas – moles reactant gas

THERMOCHEMISTRY/KINETICS

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K = -2.303 RT \log K$$

$$= -n\mathcal{F}E^\circ$$

$$\Delta G = \Delta G^\circ + RT \ln Q = \Delta G^\circ + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_p = \frac{\Delta H}{\Delta T}$$

$$\ln[\text{A}]_t - \ln[\text{A}]_0 = -kt$$

$$\frac{1}{[\text{A}]_t} - \frac{1}{[\text{A}]_0} = kt$$

$$\ln k = \frac{-E_a}{R} \left(\frac{1}{T} \right) + \ln A$$

E = energy

v = frequency

λ = wavelength

p = momentum

v = velocity

n = principal quantum number

m = mass

Speed of light, $c = 3.0 \times 10^8 \text{ m s}^{-1}$

Planck's constant, $h = 6.63 \times 10^{-34} \text{ J s}$

Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

Avogadro's number = $6.022 \times 10^{23} \text{ mol}^{-1}$

Electron charge, $e = -1.602 \times 10^{-19} \text{ coulomb}$

1 electron volt per atom = 96.5 kJ mol^{-1}

Equilibrium Constants

K_a (weak acid)

K_b (weak base)

K_w (water)

K_p (gas pressure)

K_c (molar concentrations)

S° = standard entropy

H° = standard enthalpy

G° = standard free energy

E° = standard reduction potential

T = temperature

n = moles

m = mass

q = heat

c = specific heat capacity

C_p = molar heat capacity at constant pressure

E_a = activation energy

k = rate constant

A = frequency factor

Faraday's constant, $\mathcal{F} = 96,500 \text{ coulombs per mole of electrons}$

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$

= $0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$

= $8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^\circ C + 273$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2} mv^2$$

$$KE \text{ per mole} = \frac{3}{2} RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

molarity, M = moles solute per liter solution

molality = moles solute per kilogram solvent

$$\Delta T_f = iK_f \times \text{molality}$$

$$\Delta T_b = iK_b \times \text{molality}$$

$$\pi = iMRT$$

$$A = abc$$

OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a A + b B \rightarrow c C + d D$$

$$I = \frac{q}{t}$$

$$E_{cell} = E_{cell}^\circ - \frac{RT}{nF} \ln Q = E_{cell}^\circ - \frac{0.0592}{n} \log Q @ 25^\circ C$$

$$\log K = \frac{nE^\circ}{0.0592}$$

- P = pressure
- V = volume
- T = temperature
- n = number of moles
- D = density
- m = mass
- v = velocity

u_{rms} = root-mean-square speed

KE = kinetic energy

r = rate of effusion

M = molar mass

π = osmotic pressure

i = van't Hoff factor

K_f = molal freezing-point depression constant

K_b = molal boiling-point elevation constant

A = absorbance

a = molar absorptivity

b = path length

c = concentration

Q = reaction quotient

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

E° = standard reduction potential

K = equilibrium constant

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$

$= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$

$= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

K_f for $H_2O = 1.86 \text{ K kg mol}^{-1}$

K_b for $H_2O = 0.512 \text{ K kg mol}^{-1}$

$1 \text{ atm} = 760 \text{ mm Hg}$

$= 760 \text{ torr}$

STP = $0.00^\circ C$ and 1.0 atm

Faraday's constant, $F = 96,500 \text{ coulombs per mole of electrons}$