

ANSWERS AND EXPLANATIONS TO PRACTICE EXAM I

1.

Answer: D

Topic: Acid-base

Question Type: Classification set

Question Level: Moderate

Explanation: A pH of less than 7 means an acidic buffer. This narrows the choice to C and D. Both of these choices contain an acid and a salt of the acid. However, hydrobromic acid is a strong acid and bromide ion is a very weak conjugate base. An optimum buffer is one which is composed of a weak acid and a salt of the weak acid. This allows for the buffer to resist a significant change in pH by absorbing extra H^+ and/or OH^- . Therefore, acetic acid and sodium acetate would be the only choice.

2.

Answer: E

Topic: Acid-base

Question Type: Classification set

Question Level: Moderate

Explanation: A pH greater than 7 means a basic buffer. Only choice E contains a weak base and a salt of the weak base. Choice A is a mixture of a strong base and a weak base. An optimum buffer is one which is composed of a weak base and a salt of the weak base.

3.

Answer: B

Topic: Acid-base

Question Type: Classification set

Question Level: Easy

Explanation: A pH of 7 means a neutral solution. Choice B contains a strong acid and a strong base. When mixed, the result is a neutral salt and water since the question stated that the mole to mole ratio was 1:1.

4.

Answer: A

Topic: Acid-base

Question Type: Classification set

Question Level: Moderate

Explanation: The highest pH means the most basic solution. Choice A contains a strong base and a weak base. When mixed, the pH is determined by the sodium hydroxide which is a strong base. Choice E contains only a weak base.

5.

Answer: E

Topic: Bonding

Question Type: Classification set

Question Level: Moderate

Explanation: Both MgCl_2 and NaCl are ionic compounds. The difference in melting points is due to the strength of the intramolecular forces that hold the ions together. The lattice energy or attractive force is easily estimated using Coulomb's Law, $F = \frac{q_1 q_2}{r^2}$. The difference lies in Mg's charge of +2 and Na's charge of only +1 since all of the ions are relatively the same size, the attractive force between the ions is twice as strong for MgCl_2 .

6.

Answer: E

Topic: Bonding

Question Type: Classification set

Question Level: Moderate

Explanation: Both H_2O and H_2S are covalently bonded compounds. The difference in boiling points is due to the strength of the intermolecular attractive forces that hold the molecules together. Hydrogen bonding, the attraction that hydrogen has for the unshared electron pair on the highly electronegative oxygen atom, is found in the water molecule. Hydrogen sulfide molecules are held to each other by the much weaker dipole forces. The more tightly molecules are held together, the more energy needed to separate them, and the higher the boiling point.

7.

Answer: B

Topic: Bonding

Question Type: Classification set

Question Level: Easy

Explanation: Carbon's electron configuration is $1s^2 2s^2 2p^2$. The four bonding electrons in the carbon atom are found in the s and the p orbitals. Since these orbitals are of different energy, the four carbon bonds would not be equal according to the VSEPR theory. The valence bond theory assumes that bonding electrons will hybridize to form orbitals of equal energy. In fact, in methane hybridization leads to four sp^3 hybridized orbitals.

8.

Answer: D

Topic: Bonding

Question Type: Classification set

Question Level: Easy

Explanation: The VSEPR theory relates shape of molecules to the repulsions and positions of shared and unshared electron pairs. Water obeys the octet rule and has two unshared electron pairs which together have great repulsive force on the two shared pairs which result in the bond angle being reduced from the expected 109.5° to 105° .

9.

Answer: E

Topic: Atomic Theory

Question Type: Classification set

Question Level: Moderate

Explanation: The copper +1 ion has lost one electron from the outermost energy level which is the 4th level. The resulting electron configuration for the copper +1 ion is $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^{10}$.

10.

Answer: B

Topic: Atomic Theory

Question Type: Classification set

Question Level: Easy

Explanation: The fluoride ion gains one electron to form F^{1-} . The electron configuration for the fluorine atom is $1s^2 2s^2 2p^5$. By gaining one electron, the fluoride ion becomes isoelectronic with neon, $1s^2 2s^2 2p^6$ which minimizes electron repulsion.

11.

Answer: B

Topic: Atomic Theory

Question Type: Classification set

Question Level: Easy

Explanation: Alkali metals lose their outer electron to become +1 ions. This makes all ions of the alkali metals isoelectronic with the noble gas preceding them. A sodium atom has the following electron configuration, $1s^2 2s^2 2p^6 3s^1$. By losing one electron from the highest energy level, the sodium ion becomes isoelectronic with neon which minimizes electron repulsion.

12.

Answer: A

Topic: Atomic Theory

Question Type: Classification set

Question Level: Easy

Explanation: The halogen family is group VIIA. All of the elements in this family have an “ $s^2 p^5$ ” electron configuration ending.

13.

Answer: E

Topic: Redox

Question Type: Classification set

Question Level: Moderate

Explanation: A strong oxidizing agent is one that is easily reduced. Nitric acid and potassium permanganate are both good oxidizing agents. However, the only substance listed with color is potassium permanganate. Potassium permanganate is purple in color (with its Mn^{7+}) and easily reduces to lower oxidation states which are all associated with distinct colors.

14.

Answer: A

Topic: Redox

Question Type: Classification set

Question Level: Moderate

Explanation: An oxidizing acid is one that reacts with active metals to produce oxides of that element and not the typical hydrogen gas. Nitric acid and hydrochloric acid are the two acids listed. Nitric acid, HNO_3 reacts with metals to produce nitrogen oxide compounds such as NO_2 and NO .

15.

Answer: D

Topic: Descriptive

Question Type: Classification set

Question Level: Easy

Explanation: Galvanizing is a common method of corrosion protection. Zinc is used as a coating with metals such as iron. The zinc will oxidize instead of the iron. This is very beneficial to protect the hull of ships and protect against corrosion of pipes.

16.

Answer: B

Topic: Descriptive

Question Type: Classification set

Question Level: Easy

Explanation: Acid rain is produced from gaseous nonmetal oxides (mainly SO_x and NO_x compounds, fondly referred to as “soaks and knocks”) reacting to form acids. SO_2 reacts to form acid rain in the following way. First, the sulfur dioxide combines with oxygen in the air: $2 \text{SO}_2 + \text{O}_2 \rightarrow 2 \text{SO}_3$. The sulfur trioxide combines with moisture to form the strong acid, sulfuric acid which is typically known as acid rain: $\text{SO}_3 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$

17.

Answer: B

Topic: Electrochemistry

Question Type: Classification set

Question Level: Difficult

Explanation: Adding NaCl will cause precipitation of PbCl_2 which will decrease the concentration of the lead (II) ions in solution. Using the Nernst equation, $E = E^\circ - \frac{RT}{nF} \ln Q$ where $Q = \frac{[\text{products}]^m}{[\text{reactants}]^n} = \frac{[\text{Zn}^{2+}]}{[\text{Pb}^{2+}]}$, it is easy to

see that as $[\text{Pb}^{2+}]$ decreases, Q will become greater than one and the \ln of Q will become positive and thus, the overall voltage will decrease.

18.

Answer: C

Topic: Electrochemistry

Question Type: Classification set

Question Level: Easy

Explanation: Removing the salt bridge will cause a build up of ions in each half cell, the cells will soon establish equilibrium and the voltage will drop to zero.

19.

Answer: A

Topic: Electrochemistry

Question Type: Classification set

Question Level: Difficult

Explanation: Adding water to the beaker on the left will decrease the concentration of $[Zn^{2+}]$. Using the Nernst equation, $E = E^\circ - \frac{RT}{nF} \ln Q$ where $Q = \frac{[\text{products}]^m}{[\text{reactants}]^n} = \frac{[Zn^{2+}]}{[Pb^{2+}]}$, it is easy to see that as $[Zn^{2+}]$ decreases, Q will

decrease and thus, the overall voltage will increase.

20.

Answer: B

Topic: Electrochemistry

Question Type: Classification set

Question Level: Moderate

Explanation: As current flows through a cell, the concentration of the ions being oxidized and reduced are constantly decreasing. The voltage of the cell will continue to decrease over time until it reaches equilibrium, zero.

21.

Answer: B

Topic: Laboratory

Question Type: Multiple choice

Question Level: Moderate

Explanation: Solubility rules must be memorized! KI is a soluble salt and forms K^+ and I^- ions in solution. The ions of alkali metals do not form precipitates so this eliminates choices C, D, and E. Iodide ions are soluble with everything except for Ag^+ , Pb^{2+} and Hg_2^{2+} . The net ionic reaction is: $Pb^{2+} + 2 I^- \rightarrow PbI_2$

22.

Answer: C

Topic: Equilibrium

Question Type: Multiple choice

Question Level: Moderate

Explanation: The negative value for ΔH given means that the reaction is exothermic (energy is given off as a product). If the temperature is decreased, the reaction will shift to the product side to replace the heat. Pressure will have no effect since the same number of moles of gas is present on either side of the equation. Adding an inert gas or adding a catalyst has no effect on the equilibrium position since neither appears in the equilibrium expression since they are neither a reactant nor a product. Allowing hydrogen to escape would decrease the concentration of a reactant and thus, shift the equilibrium to the reactant side.

23.

Answer: C

Topic: Acid-Base

Question Type: Except

Question Level: Moderate

Explanation: A Brønsted acid donates a proton while a Brønsted base accepts a proton. The following possibilities exist:

<i>Substance</i>	<i>Acid</i>	<i>Base</i>
HSO_3^-	H_2SO_3	SO_3^{2-}
HPO_4^{2-}	H_2PO_4^-	PO_4^{3-}
NH_4^+	NH_3	NH_5^{2+}
H_2O	H_3O^+	OH^-
HCO_3^-	H_2CO_3	CO_3^{2-}

Nitrogen is not capable of expanding its octet to accommodate five hydrogen atoms nor is NH_3 an acid.

24.

Answer: D

Topic: Laboratory

Question Type: Multiple choice

Question Level: Moderate

Explanation: The only significant discrepancy in the data is between the first and subsequent samples. In a titration, the number of moles of acid must equal the number of moles of base at the equivalence point. Adding more water would just increase the volume of the acid but would not affect the number of moles of acid.

Adding more phenolphthalein has no effect. If the sample was titrated beyond the endpoint or the buret not rinsed with NaOH the volume of NaOH would appear too large. By not rinsing the pipet with acetic acid, the error is introduced.

25.

Answer: B

Topic: Thermodynamics

Question Type: Multiple choice

Question Level: Easy

Explanation: The most positive value of ΔS indicates the system with the most disorder as the reaction proceeds from reactants to products. Choice B shows 2 moles of gaseous reactant producing 3 moles of gaseous product. All other choices show the formation of solids and/or liquids which is a decrease in entropy.

26.

Answer: A

Topic: Periodicity

Question Type: Multiple choice

Question Level: Easy

Explanation: Moving across a period from Li to Ne, electrons are being added to the same energy level.

However, the number of protons is also increasing which causes a greater effective nuclear charge which allows for the outer electrons to be held more tightly, thus, decreasing the size.

27.

Answer: A

Topic: Gases

Question Type: Multiple choice

Question Level: Moderate

Explanation: $14 \text{ g N}_2 \times \frac{1 \text{ mol}}{28 \text{ g N}_2} = .50 \text{ mol N}_2$ $22 \text{ g CO}_2 \times \frac{1 \text{ mol}}{44 \text{ g CO}_2} = .50 \text{ mol CO}_2$

Since both compounds contain the same number of moles at the same temperature and pressure the volume occupied by each would be the same as would the number of molecules. The density of a gas is reported as grams per liter; nitrogen gas is less dense than carbon dioxide. Average kinetic energy is temperature dependent and all molecules at the same temperature have the same average kinetic energy. Average speed is inversely proportional to the square root of the molecular weight. In general, lighter molecules will travel faster than heavier molecules so nitrogen would move more quickly than carbon dioxide.

28.

Answer: E

Topic: Bonding

Question Type: Multiple choice

Question Level: Easy

Explanation: The question states that XeF₄ is octahedral, which means it has 6 sites of electron density. That means the hybridization uses the s orbital plus all three p orbitals, plus two d orbitals for a total of six “sites” of electron density. The hybridization of atomic orbitals creates bonding areas or “sites” of equal energy. The following hybridizations are common:

<i>Hybridization</i>	<i>Electronic Shape</i>
sp	linear
sp ²	trigonal planar
sp ³	tetrahedral
dsp ³	trigonal bipyramidal
d ² sp ³	octahedral

29.

Answer: B

Topic: Kinetics

Question Type: Multiple choice

Question Level: Moderate

Explanation: In experiment #1 and #2 [B] is held constant while [A] doubles and the rate doubles. The reaction is 1st order with respect to A. In experiment #2 and #3 [A] is held constant while [B] is increased 5 times and the rate increases 25 times. Setting this up mathematically makes it easy to see that the reaction is 2nd order with respect to B.

$5^n = 25$ so ... $n = 2$ (or second order).

30.

Answer: C

Topic: Acid-Base

Question Type: Multiple choice

Question Level: Easy

Explanation: In the experiment the flask begins with a weak base. A base has a pH greater than 7 so identifying this narrows the choice to C, D, and E. A weak base typically has a pH less than 12 thus eliminating D and E.

31.

Answer: B

Topic: Nuclear

Question Type: Multiple choice

Question Level: Easy

Explanation: In transmutation reactions, the superscripts and the subscripts on each side must be equal. Total superscripts on the reactant side equals 247. On the product side the coefficient of two multiplies to give a total of 2 (1 for each of two neutrons) + 129 (for iodine) = 131. Subtract: $247 - 131 = 116$ which represents the top number. This narrows the choice to A or B. Follow the same procedure for the subscript. Total on reactant side = 97. Total on product side = 53. Subtract: $97 - 53 = 44$. Ruthenium-44 is the correct choice.

32.


Answer: C

Topic: Bonding

Question Type: Multiple choice

Question Level: Easy

Explanation: Sketch each Lewis structure to determine number of unshared electron pairs on the central atom.



Bromine and oxygen have no central atom. Methane, CH_4 , has a molecular geometry that is tetrahedral and has no unshared electron pairs. SO_3 has a molecular geometry that is trigonal planar with no unshared electron pairs on the central atom while ammonia has a molecular geometry that is trigonal pyramidal and has one unshared pair of electrons on the central nitrogen atom.

33.


Answer: C

Topic: Kinetics


Question Type: Multiple choice

Question Level: Moderate


Explanation: Determining the order from experimental data requires graphing. If create the set of graphs in this order with the y-axes being “concentration”, “natural log of concentration” and “reciprocal concentration” (Alphabetical order by axes names), whichever gives a straight line corresponds to zero, first, or second order. (Alphabetical order of the y-axis variables \rightarrow 0, 1, 2 orders for that reactant). Time is always on the x-axis.



Zero order
 $k = \text{negative slope.}$



First order
 $k = \text{negative slope.}$



Second order
 $k = \text{the slope.}$

34.

Answer: C

Topic: Thermodynamics

Question Type: Multiple choice

Question Level: Easy

Explanation: If the reaction is spontaneous, ΔG is negative and the second condition of spontaneity is that K_{eq} be greater than one.

35.


Answer: E

Topic: Bonding

Question Type: Multiple choice

Question Level: Difficult

Explanation:



Pi bonds are present when multiple bonds are present. The structures for each of the choices must be drawn. Chloric acid, HClO₃, is the only compound with all single bonds and is easily eliminated. Acetic acid and sulfur dioxide each have one pi bond and carbon dioxide has two pi bonds. Benzene, C₆H₆, is a ring structure with three double bonds, thus, the most pi bonding.

36.

Answer: C

Topic: Descriptive

Question Type: Multiple choice

Question Level: Easy

Explanation: Potassium, manganese, and uranium are all solid metals at room temperature while chlorine is a yellow-green gas. Bromine is the only liquid in the list.

37.

Answer: A

Topic: Descriptive

Question Type: Multiple choice

Question Level: Moderate

Explanation: Alkali metals react violently with water to produce strong bases. The reaction is:



38.

Answer: E

Topic: Redox

Question Type: Multiple choice

Question Level: Moderate

Explanation: Assign oxidation numbers to each element in the reaction then analyze. Oxygen changes from 0 to -2. H stays +1. Ag changes from 0 to +1. C is +2 in the cyanide ion. N is -3 in the cyanide ion. Only two elements may change oxidation state within one reaction; one must be oxidized and one must be reduced. If carbon were +4 in the cyanide ion, the charge on the ion would be +1 instead of -1.

39.

Answer: B

Topic: Electrochemistry

Question Type: Multiple choice

Question Level: Moderate

Explanation: The half reaction given is written as a reduction since the electrons are on the reactant side. Analyzing oxidation numbers, it is easy to see that gold changes from +3 to 0. This represents a reduction. Reduction always occurs at the cathode. The chloride ion does not change charge in this half reaction. The reducing agent is not shown here since it is the substance undergoing oxidation.

40.

Answer: C

Topic: Electrochemistry

Question Type: Multiple choice

Question Level: Moderate

Explanation: The choices give the numerical set up without units. Write the units in and then see what matches numerically. Remember that an amp is equal to a coulomb per second.

$$\frac{2.0 \text{ coulomb}}{\text{s}} \times \frac{10 \text{ min}}{1} \times \frac{60 \text{ s}}{1 \text{ min}} \times \frac{1 \text{ mol e}^-}{96,500 \text{ coulombs}} \times \frac{1 \text{ mol Au}}{3 \text{ mol e}^-} \times \frac{197 \text{ g Au}}{1 \text{ mol Au}} =$$

41.

Answer: E

Topic: Nuclear

Question Type: Multiple choice

Question Level: Moderate

Explanation: Half-life is the time that it takes for $\frac{1}{2}$ of the substance to decay. Without a calculator, it is best to set this problem up and work backwards for every 5 minutes since this is the $\frac{1}{2}$ life given in the problem.

<i>Amount</i>	<i>Time(min.)</i>
80 g	45
160 g	40
320 g	35
640 g	30
1280g	25
2560 g	20
5120 g	15
10,240 g	10
20,480 g	5
40,960 g	0 (this is original amount at the beginning)

42.

Answer: B

Topic: Atomic theory

Question Type: Multiple choice

Question Level: Easy

Explanation: Rutherford expected that all of the positively-charged alpha particles would be attracted to the sea of electrons and pass straight through the foil. To his surprise, some of the positive particles were deflected and bounced back. This meant that there must be some dense, positive substance within the atom. Many of the answer choices given in this question are true statements but are not significant to Rutherford's experiment.

43.

Answer: D


Topic: Thermodynamics

Question Type: Multiple choice

Question Level: Difficult

Explanation: The desired equation is: $6\text{C}(\text{s}) + 3\text{H}_2(\text{g}) \rightarrow \text{C}_6\text{H}_6(\text{l})$


The three equations given must be rearranged to give this equation. Remember that for thermodynamic data, whatever is done to the equation is done to the thermodynamic value. If the equation is reversed, change the sign. If the equation is multiplied by some value, the thermodynamic quantity is multiplied by that same value. The first equation is reversed and divided by 2:



The second equation is multiplied by 6:



The third equation is multiplied by 3:



Adding the equations together gives the desired reaction. Add the ΔG° values together $3200. \text{ kJ} + (-3150. \text{ kJ}) = 50. \text{ kJ}$

44.

Answer: A

Topic: Kinetics

Question Type: Multiple choice

Question Level: Easy

Explanation: If $[\text{Y}]$ doubles and the rate quadruples all other factors held constant then the reaction must be 2nd order with respect to $[\text{Y}]$. $2^n = 4$; $n = 2$. We do not have enough information about the reaction mechanism to determine choices B, C, or D. Choice E is an incorrect statement.

45.

Answer: E

Topic: Laboratory

Question Type: I, II, III

Question Level: Easy

Explanation: All of the statements are appropriate lab procedures. pH probes must be calibrated in order to give accurate readings. Glass tubing slides easily into a rubber stopper with lubrication but may break if lubrication is not used. Massing an object that is hot will cause the measured mass to be less than the true value because of the convection currents produced by when hot air collides with the cold air around the dish.

46.

Answer: A

Topic: Kinetics

Question Type: Except

Question Level: Easy

Explanation: Catalysts generally speed up reaction rate by providing alternate pathways which take less energy for reactants to become products. All of the other choices given slow reaction rates down. By lowering the temperature, molecules collide less often with less force. Increasing the concentration of a product may shift the reaction to favor the reactants. High bond energy and high activation energy both refer to a large amount of energy needed to overcome the activation energy barrier which is indicative of slower reactions.

47.


Answer: D

Topic: Stoichiometry

Question Type: Multiple choice



Question Level: Moderate

Explanation: The net ionic reaction for the precipitation of barium with sulfate is:



The total moles of barium in the solution will be equal to the total moles of sodium sulfate needed for complete precipitation.

Find the total moles of barium in the solution:



Total Ba^{2+} in 1.0 liter of solution is: $0.10 \text{ mol} + 0.30 \text{ mol} = 0.40 \text{ moles}$. This is the number of moles of Na_2SO_4 needed.

48.

Answer: A

Topic: Equilibrium

Question Type: Multiple choice

Question Level: Moderate

Explanation: Products are favored at equilibrium since the given K value is greater than one. Ammonia, NH_3 , and acetate ion, $\text{C}_2\text{H}_3\text{O}_2^-$, both act as bases but since the reaction proceeds as written, ammonia must be a stronger base. Using the same logic, choice B is incorrect since acetic acid, $\text{HC}_2\text{H}_3\text{O}_2$, must be the stronger acid since the reaction proceeds in the forward direction. Choice C is incorrect since acetate ion is the conjugate base for acetic acid and not for ammonia. There is not enough information given to know whether choice D and E are correct.

49.

Answer: C

Topic: Equilibrium

Question Type: Multiple choice

Question Level: Difficult

Explanation: Given the equation: $2 A(g) + B(g) \rightleftharpoons 2 C(g)$ the K_c expression is: $K_c = \frac{[C]^2}{[A]^2[B]}$

Using the information provided and the stoichiometry calculate equilibrium concentrations.

2 A	+	B	\rightleftharpoons	2 C
.60 mol		.75 mol		0
-2x		-x		+2x
?		?		.30 mol

Knowing $2x = .30$ then $x = .15$ calculate the concentrations of A and B at equilibrium.

$$[A] = .60 \text{ mol} - .30 \text{ mol} = .30 \text{ mol}$$

$$[B] = .75 \text{ mol} - .15 \text{ mol} = .60 \text{ mol}$$

The volume is 1.0 liter so moles = molarity in this problem. Plug in values and estimate the answer.

$$K_c = \frac{[.30]^2}{[.30]^2 [.60]}$$

The problem simplifies to 1 divided by .60, which is the reciprocal of

$$\frac{6}{10} \text{ so the correct answer is } 10 \div 6 \approx 1.7.$$

50.


Answer: C

Topic: Bonding

Question Type: Multiple choice

Question Level: Moderate

Explanation:



Draw each structure listed and determine their shapes.

Molecule	Shape
CH_4	tetrahedral
SiO_2	linear
$AsCl_3$	trigonal pyramidal
$AsCl_5$	trigonal bipyramidal
SF_6	octahedral

A square planar molecule has 4 shared pairs of electrons all in one plane and one unshared pair of electrons above and below the plane.

51.

Answer: B

Topic: Acid-Base

Question Type: Multiple choice

Question Level: Difficult

Explanation: $\text{pH} = -\log [\text{H}^+]$ and $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$

Plugging in values makes the estimation a bit easier. $6.5 \times 10^{-10} = \frac{[x][x]}{[1.0 \times 10^{-3}]}$ Cross multiplying gives $6.5 \times$

10^{-13} ; the square root of the exponent would be around -6.5. Log base 10 is the exponent so the only choice in this range of 6 is choice B.

52.

Answer: D

Topic: Solutions

Question Type: Multiple choice

Question Level: Moderate

Explanation: This appears to be a simple dilution problem. $M_1V_1 = M_2V_2$ Plugging in values leads to: $(6.00 \text{ M})(10.0 \text{ mL}) = (0.100 \text{ M})(V_2)$

$V_2 = 600.0 \text{ mL}$ which represents the total volume of the solution. The trick here is that the problem asks for how much water should be added to the 10.0 mL of acid. Subtracting: $600.0 \text{ mL} - 10.0 \text{ mL} = 590. \text{ mL}$ of water needed.

53.

Answer: C

Topic: Laboratory

Question Type: I, II, III

Question Level: Moderate

Explanation: A visible-light spectrophotometer is useful for determining the concentration of colored solutions by measuring either the % transmittance of light passing through a solution or by measuring the absorbance of light by the solution. The absorbance is directly proportional to the concentration according to Beer's law; $A = \epsilon bc$. All zinc solutions are colorless since all of its d electrons are paired, so choice I is eliminated. Potassium permanganate is colored but conductivity is not measured by light passing through the substance so choice II is eliminated. All of the ions listed in the third choice are colored in solution so determining their concentrations would be possible using a spectrophotometer.

54.

Answer: E

Topic: Acid-Base

Question Type: Multiple choice

Question Level: Difficult

Explanation: A Lewis acid is an electron pair acceptor. Fluoride ion and perchlorate ion are both negative and therefore, have no affinity for accepting an electron pair. The choice is narrowed to the metal ions listed. The metal with the highest positive charge has the highest affinity for an electron pair, which in this case is the aluminum ion.

55.

Answer: E

Topic: Electrochemistry

Question Type: Multiple choice

Question Level: Moderate

Explanation: Given that the K_{eq} value is less than one indicates that the reactants are favored at equilibrium. This indicates that the reaction proceeds in the reverse direction. The voltage of the cell would be negative and the ΔG value would be positive.

56.

Answer: B

Topic: Laboratory

Question Type: Multiple choice

Question Level: Easy

Explanation: The net ionic reaction is: $SCN^- + Fe^{3+} \leftrightarrow FeSCN^{2+}$. This complex ion is known to have a distinct brick red color. A bright yellow color might indicate a precipitate of lead (II) iodide. A solution changing from light blue to dark blue might indicate a complex of copper. Bubbles of nitrogen dioxide usually form from a metal reacting with concentrated nitric acid and the pH decreasing indicates hydrogen ions being produced in solution.

57.

Answer: D


Topic: Electrochemistry

Question Type: Multiple choice


Question Level: Difficult

Explanation: The desired half-reaction is: $X^{3+}(aq) + 3 e^- \rightarrow X(s)$ It must be obtained from the two given equations. The equations may be reversed, multiplied and/or divided before summing them to obtain the desired equation. If the reaction is reversed, the sign on the voltage will reverse. The trick here is that voltages are not a "per mole" quantity and thus, are not multiplied or divided by anything.


Equation one is reversed:



Equation two is multiplied by 3:



Adding these two equations gives:



This equation needs to be divided by two to look like the desired equation, but the voltage will just be the addition of the reactions above. $E = 3.87 + (-2.37) = 1.50$ volts

58.

Answer: E

Topic: Laboratory

Question Type: Multiple choice

Question Level: Easy

Explanation: The easiest way to recover sodium chloride is to evaporate to dryness. Distillation is a separation method for liquids with different boiling points. Filtration works well when particles are suspended in a solution. Electrolysis is useful for obtaining pure forms of elements by running an electric current through either the molten state or a solution. Fractional crystallization is a method for separating multiple solutes contained in a solution. The solution is heated and the various solutes crystallize at different temperatures as the solution cools.

59.

Answer: D

Topic: Laboratory

Question Type: I, II, III

Question Level: Difficult

Explanation: There are two concepts to keep in mind. First, in any titration moles acid = moles base at the equivalence point. Secondly, K_a is the acid dissociation constant. Failure to rinse the buret with the NaOH after washing, means that water droplets adhering to the walls of the buret will dilute the NaOH solution so more base will be required and the moles of base reported will be too high so the moles of acid, HA, reported will also be too high implying more HA dissociated, so that the acid dissociation constant, K_a , is reported as too large. Not adding enough water to dissolve the entire solid acid sample will lower the number of acid molecules that dissociate in solution so less base will be required, and the moles of acid and thus the acid dissociation constant will be reported as too small. The addition of base beyond the equivalence point causes the moles of base reported to be too large, thus the moles of acid reported are also too large, so the acid dissociation constant is reported as too large.

60.

Answer: B

Topic: Periodicity

Question Type: Multiple choice

Question Level: Easy

Explanation: The largest jump in ionization energy is between removal electron number 2 and electron number 3. Large incremental increases in IE usually indicate the removal of an electron from an inner energy level.

Write electron configurations for the choices.

<i>Element</i>	<i>Electron Configuration</i>
K	[Ar] 4s ¹
Ca	[Ar] 4s ²
Ga	[Ar] 4s ² 3d ¹⁰ 4p ¹
Ge	[Ar] 4s ² 3d ¹⁰ 4p ²
As	[Ar] 4s ² 3d ¹⁰ 4p ³

The largest increase in IE for K would be between the removal of electron #1 and #2. The largest increase in IE for Ca is between electron #2 and #3 since the previous principle energy level is broken into. For Ga, Ge, and As the largest jumps would occur between #3 and #4; #4 and #5; #5 and #6 respectively.

61.


Answer: B

Topic: Stoichiometry

Question Type: Multiple choice

Question Level: Moderate

Explanation: Barium hydroxide is a strong base and would ionize close to 100% according the following equation:



There would be relatively zero $\text{Ba}(\text{OH})_2$ present in a 0.500 M solution. The concentration of Ba^{2+} ions would be 0.500 M since the ratio is 1:1 with $\text{Ba}(\text{OH})_2$. The concentration of OH^- is $2(.500) = 1.00$ M since the ratio is 1:2 with $\text{Ba}(\text{OH})_2$. The substance BaOH^+ does not exist. Hydronium ions, H_3O^+ ions would not exist in a strong basic solution.

62.

Answer: C

Topic: Solutions

Question Type: Multiple choice

Question Level: Difficult

Explanation: $MM = \frac{\text{grams}}{\text{moles}}$ Given this equation, we must know the mass in grams of the substance and the

number of moles of the substance. Given the information about the difference in temperature and the boiling point constant, molality could be found using the following equation. $\Delta T_b = K_b \cdot m \cdot i$ Since the unknown substance is molecular and nonvolatile the i value is 1. Using the molality formula, moles could be calculated if the mass of solvent is known. $m = \frac{\text{mol solute}}{\text{kg solvent}}$ So...in addition to the change in boiling point and the boiling

point constant, the mass of unknown (solute) used and the mass of solvent used would be necessary to determine the molecular mass.

63.

Answer: C

Topic: Stoichiometry

Question Type: Multiple choice

Question Level: Moderate

Explanation: A hydrocarbon is composed of hydrogen and carbon. Assume a 100% sample, or 100. gram sample and simplify the problem. If 20.0%, or 20.0 grams, is due to hydrogen then 80.0%, or 80.0 grams, is due to carbon. First convert the grams into moles for each substance.

$$20.0 \text{ g H} \times \frac{1 \text{ mol H}}{1.01 \text{ g H}} = 19.88 \text{ mol H}$$

$$80.0 \text{ g C} \times \frac{1 \text{ mol C}}{12.01 \text{ g C}} = 6.66 \text{ mol C}$$

Next, divide each of the moles by the smaller number of moles to find subscripts. $6.66/6.66 = 1$; $19.88/6.66$ is about 3. The formula is CH_3 .

64.


Answer: D

Topic: Reaction types

Question Type: Multiple choice

Question Level: Difficult

Explanation: A solution saturated with silver chloride would appear cloudy and the precipitate would be white. Silver chloride is not very soluble and is thus represented as an insoluble salt. Bubbling ammonia gas into the solution would clear the solution as a soluble complex ion forms. The balanced net ionic reaction would be:



65.

Answer: E

Topic: Solutions

Question Type: Multiple choice

Question Level: Easy

Explanation: Choices A, B, and C are all soluble salts and form ions in solution. Choice B is a weak acid and will produce some ions in solution. However, ethanol, $\text{C}_2\text{H}_5\text{OH}$ is an alcohol. Organic alcohols do not ionize in solution. Be careful not to mistake the alcohol for a base!

66.


Answer: B

Topic: Acid-Base

Question Type: Multiple choice

Question Level: Moderate

Explanation: Oxidation of an oxyacid means the addition of more oxygen to the acid as well as change in oxidation number. The more oxygen the acid contains in a series of acids, the stronger the acid becomes due to the highly electronegative oxygen atoms pulling on the electrons polarizing the O-H bond. (It is also important to note that the hydrogen that is to be ionized (removed) is bound to one of the oxygen atoms of the oxyacid.) When HNO_2 , a weak acid, is oxidized to HNO_3 , a strong acid, the oxidation number of nitrogen changes from +3 to +5 which indicates a loss of electrons or oxidation.



67.

Answer: E

Topic: Stoichiometry

Question Type: Multiple choice

Question Level: Moderate

Explanation: This is a limiting reactant problem. Write a balanced equation first. Notice that everything is in a 1:1 ratio.

Ba(NO ₃) ₂	+	MgSO ₄	→	Mg(NO ₃) ₂	+	BaSO ₄
100. mL		100. mL				
0.10 M		0.060 M				
10.0 mmol		6.00 mmol	**MgSO ₄ limits the reaction			
-6.00mmol		-6.00 mmol				
4.00 mmol left		0.00 mmol				

Calculate the number of moles (or millimoles) of given reactants to determine which one is limiting. Just multiply molarity x volume to get millimoles. Without a calculator this is easy to do since you are just moving decimals. Determine the limiting reactant—in this case since everything is 1:1 the smaller number limits. Subtract the limiting reactant from the excess reactant to see how many moles are left. Calculate the new molarity of Ba²⁺ ions left by dividing mmoles by total volume.

$$M = \frac{\text{mmol Ba}^{2+}}{\text{total volume solution}}$$

$$M = \frac{4.00 \text{ mmol Ba}^{2+}}{200 \text{ mL}} = 0.02$$

68.

Answer: E

Topic: Liquids/solids

Question Type: Multiple choice

Question Level: Moderate

Explanation: In the diagram below point A to C represents the liquid phase; Points B to C represent supercooling of the liquid; Points C to D represent melting and freezing; Points D to E represent the solid phase.

69.

Answer: E

Topic: Gases

Question Type: Multiple choice

Question Level: Moderate

Explanation: In this gas law problem both number of moles and volume are constant. This makes a simple Gay-Lussac law problem. The formula is: $\frac{P_1}{T_1} = \frac{P_2}{T_2}$ The temperatures must be converted to Kelvin before

solving. The equation becomes: $\frac{2.5 \text{ atm}}{400 \text{ K}} = \frac{P_2}{300 \text{ K}}$ To solve, cross-multiply and simplify. $750 / 400$ is close to the number 2. There is only one answer close to this—1.88 atm.

70.

Answer: E

Topic: Solids/Liquids/Gases

Question Type: Multiple choice

Question Level: Moderate

Explanation: “Normal” boiling point is the point at which the liquid and vapor are in equilibrium at standard atmospheric pressure. In the phase diagram given, when the substance is at 1 atm of pressure, the liquid-vapor equilibrium line does not cross the dashed line of 1atm so the boiling point is undetermined by this graph.

71.

Answer: B

Topic: Solids/Liquids/Gases

Question Type: Multiple choice

Question Level: Easy

Explanation: Only the conditions necessary for sublimation, the change from gas to solid, are provided by the phase diagram. Density determination requires data of mass and volume. Specific heat, latent heat of vaporization and latent heat of fusion are given by typical heating curves not phase diagrams.

72.

Answer: A

Topic: Solids/Liquids/Gases

Question Type: Multiple choice

Question Level: Moderate

Explanation: The gaseous state of all substances is less dense than either the solid or the liquid phase of the same material because there are very few particles of gas in a given volume. The slope of the solid-liquid equilibrium line gives qualitative data about the density of the solid in comparison to the liquid. In this example, the slope of the solid-liquid equilibrium line is positive. This is most common and means that the solid is more dense than the liquid.

73.

Answer: B

Topic: Stoichiometry

Question Type: Multiple choice

Question Level: Easy

Explanation: The first step in solving a stoichiometric problem is to write a balanced equation. $\text{Al}_2\text{S}_3 + 3 \text{H}_2 \rightarrow 2 \text{Al} + 3 \text{H}_2\text{S}$ Notice the ratio between aluminum sulfide and aluminum is 1:2. This makes the math

easy

$$.500 \text{ mol Al}_2\text{S}_3 \times \frac{2 \text{ mol Al}}{1 \text{ mol Al}_2\text{S}_3} \times \frac{27.0 \text{ g Al}}{1 \text{ mol Al}} = 27.0 \text{ g Al}$$

74.

Answer: B

Topic: Stoichiometry

Question Type: Multiple choice

Question Level: Moderate

Explanation: $\% = \frac{\text{mass of MgCO}_3}{\text{total mass of compound}} \times 100$ The only thing we begin the problem knowing is that the total

mass of the compound is 16.8 grams. We must write a balanced equation to see how CO_2 and MgCO_3 are related. $\text{MgCO}_3 \rightarrow \text{MgO} + \text{CO}_2$ There is a 1:1 relationship between carbon dioxide and magnesium carbonate. The set up might look something like this—

$$4.4 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44 \text{ g CO}_2} \times \frac{1 \text{ mol MgCO}_3}{1 \text{ mol CO}_2} \times \frac{84 \text{ g MgCO}_3}{1 \text{ mol MgCO}_3} = 8.4 \text{ g MgCO}_3$$

Now, plug the mass of magnesium carbonate into the first formula to find the percent by mass.

$$\% = \frac{8.4 \text{ g MgCO}_3}{16.8 \text{ g total}} \times 100 = 50\%$$

75.

Answer: A

Topic: Redox

Question Type: Multiple choice

Question Level: Easy

Explanation: Given that the two reactants are in 1:1 ratio with each other, the number of Cr atoms in the product must be 2. $..1.. \text{Cr} + ..1.. \text{CrO}_4^{2-} + \dots \rightarrow \dots 2 \text{Cr(OH)}_3$